

**Study Phase Plan: MutaFlow<sup>BASIC</sup> Pig-a Analysis Kit (P15MFv16)  
Frozen Rodent Blood**

An original signed Study Phase Plan document and Sample Submission Form are required for each study. If requesting GLP analysis, a copy of your protocol is also required prior to sample analysis.

**A. Contact Information**

Test Facility Name and Address:

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Test Facility Study Director:

Name \_\_\_\_\_

Phone \_\_\_\_\_

Fax \_\_\_\_\_

Email \_\_\_\_\_

Test Site Name and Address:

Litron Laboratories  
3500 Winton Place, Suite 1B  
Rochester, New York 14623  
phone: 585-442-0930  
fax: 585-442-0934  
info@litronlabs.com  
www.litronlabs.com

**B. Study Information**

Study ID: \_\_\_\_\_ Indicate Species: \_\_\_\_\_ Rat \_\_\_\_\_ Mouse Indicate Strain: \_\_\_\_\_

For analysis in compliance with GLP regulations, initial here. Provide the agency name that the data will be submitted to: \_\_\_\_\_. Also, indicate which GLP regulations should be followed (FDA and/or OECD): \_\_\_\_\_. A copy of your protocol is required. For FDA GLP analyses, label samples with Sample ID, Study ID, Date Collected, Source (i.e., mouse or rat) and Type (i.e., blood). For OECD GLP, label samples with Unique ID and Sample ID.

For Non-GLP analysis, initial here if a study phase report is requested in addition to the electronic data file.

Initial here for statistical analysis of data. Additional charges apply. Contact Litron for details.

All study phase specific records will be sent to your test facility after study phase completion, otherwise initial here to have records retained at Litron (see Section 8).

**If applicable, please indicate any requested modifications to the Study Phase Plan:**

\_\_\_\_\_

**C. Study Phase Plan Approval**

**Study Director Signature:** \_\_\_\_\_ **Date:** \_\_\_\_\_

**For Litron use only**

Principal Investigator \_\_\_\_\_ GLP Number: \_\_\_\_\_

Principal Investigator's Signature \_\_\_\_\_ Date \_\_\_\_\_

## 1. Objective

This MutaFlow® Study Phase Plan describes procedures for analyzing test facility-submitted frozen mouse or rat blood samples and determining the frequencies of mutant phenotype erythrocytes (RBCs), mutant phenotype immature erythrocytes (reticulocytes, or RETs), and RETs using flow cytometry. The method is based on the endogenous *Pig-a* gene whose product is essential for the synthesis of glycosylphosphatidylinositol (GPI) anchors.

## 2. Introduction

Hematopoietic cells require GPI anchors to attach a host of proteins to their cell surface, for instance, CD24, CD59, and CD55. Importantly, of the genes required to form GPI anchors, only *Pig-a* is located on the X-chromosome. Mutations in the *Pig-a* gene can prevent functional anchors from being produced, resulting in cells lacking these proteins on their surface. Thus, cells without these cell surface markers represent a reliable phenotypic marker of *Pig-a* mutation. Flow cytometry is used for this analysis, as it provides a high-speed method for objective scoring of these very rare events. *Pig-a* mutation occurs spontaneously at a very low frequency, but mutagens cause an increase in the number of mutants relative to the background (spontaneous) level.

## 3. Proposed Study Dates

The experimental start and end dates will be documented in the raw data.

## 4. Experimental Procedures (performed at Test Facility)

The test facility is responsible for following the procedures detailed in the Litron-provided manual. Deviations from the procedures described in the manual are not recommended. Modifications not previously approved by Litron may result in samples that are incompatible with flow cytometric analysis. In addition to this Study Phase Plan, a Sample Submission Form should accompany each shipment of samples to Litron.

## 5. Flow Cytometric Analysis (performed at Test Site)

### • Sample Receipt, Labeling and Column Fractionation

Upon receipt at Litron, frozen blood samples will be either immediately thawed or stored in a -75 °C to -85 °C freezer and thawed at a later date. After thawing, samples are processed through Lympholyte®-Mammal Solution to remove the majority of leukocytes and platelets.

Cells are then incubated with Anti-CD24-PE (mouse) or Anti-CD59-PE (rat) to label wild-type (wt) erythrocytes and Anti-CD61-PE (to label remaining platelets). Antibody-labeled samples are incubated with Anti-PE MicroBeads, which bind to these antibodies. A small fraction of each sample is stained with a nucleic acid dye (to differentiate leukocytes and RETs from mature RBCs). This dye solution also includes fluorescent Counting Beads and these “Pre-Column” samples are analyzed to capture Cell:Bead ratios.

The remaining portion (majority) of the blood sample is applied to a column that has been suspended in a magnetic field. These columns selectively retain wt cells, whereas *Pig-a* mutant phenotype cells (lacking CD24 or CD59 on their surface) pass through the columns.

Eluates are collected and centrifuged, stained with a nucleic acid dye (to differentiate leukocytes and RETs from mature RBCs). This dye solution also includes fluorescent Counting Beads and these “Post-Column” samples are analyzed on a flow cytometer to capture Mutant Cell:Bead ratios.

### • Flow Cytometer Calibration

An Instrument Calibration Standard sample will be used to configure and calibrate the flow cytometer before analysis.

### • Analysis of Samples

Samples will be analyzed by flow cytometry. The stained cells are moved past a laser set to provide 488 nm excitation. The fluorescence emitted by each cell is collected by photomultiplier tubes.

## 6. Data Provided

For Pre-Column samples, a stop mode will be used based on the length of time needed to acquire at least 1,000 Counting Beads.

For Post-Column samples, a stop mode will be used based on the length of time needed to analyze nearly the entire volume of cells and Counting Beads.

From the Pre- and Post-Column analyses, the following values are calculated:

- Frequency of RETs, an index of bone marrow toxicity, expressed as percent of total RBCs
- Frequency of mutant-phenotype RBCs, expressed as number per 1,000,000 total RBCs
  - Note that upon acute mutagen exposure this index of genotoxicity is not expected to reach a maximal response until the entire cohort of circulating RBCs has turned over (approximately 30 days for mice, and approximately 60 days for rats).
- Frequency of mutant-phenotype RETs, expressed as number per 1,000,000 total RETs
  - Note that upon acute mutagen exposure this index of genotoxicity reaches a maximal value faster than mutant phenotype RBCs (often within approximately 2 weeks), since RETs are turned over at a much faster rate than the total RBC pool.

## 7. Evaluation and Interpretation of Results

No statistical analyses will be performed on the data, other than the calculations indicated above, and the test facility will be responsible for the evaluation and interpretation of results.

## 8. Records Maintained

The original study phase plan and study-specific records will be transferred to the test facility at the completion of the study phase. Litron will maintain a copy of the study phase plan and copies of the paper and electronic records for two years following completion of the analysis. After the retention period, Litron will contact the sponsor and these items will either be discarded or sent to the sponsor-requested facility.

## 9. References

- Avlasevich SL, Phonetheswath S, Labash C, Carlson K, Torous DK, Cottom J, Bemis JC, MacGregor JT, and Dertinger SD (2014) *Environmental & Molecular Mutagenesis* 55, 400-406.
- Bhalli JA, Ding W, Shaddock JG, Pearce MG, Dobrovolsky VN, and Heflich RH (2013) *Mutation Research* 753, 82-92.
- Bhalli JA, Shaddock JG, Pearce MG, Dobrovolsky VN, Mutagenesis (2013) 28, 447-455.
- Bhalli JA, Shaddock JG, Pearce MG, Dobrovolsky VN, Cao X, Heflich RH, and Vohr HW (2011) *Environmental & Molecular Mutagenesis* 52, 731-737.
- Cammerer Z, Bhalli JA, Cao X, Coffing SL, Dickinson D, Dobo KL, Dobrovolsky VN, Engel M, Fiedler RD, Gunther WC, Heflich RH, Pearce MG, Shaddock JG, Shutsky T, Thiffeault CJ, and Schuler M (2011) *Environmental & Molecular Mutagenesis* 52, 721-730.
- Dertinger SD, Avlasevich SL, Torous DK, Bemis JC, Phonetheswath S, Labash C, Carlson K, Mereness J, Cottom J, Palis J, and MacGregor JT (2014) *Toxicological Sciences* 140, 307-314.
- Dertinger SD, Bryce SM, Phonetheswath S, Avlasevich SL (2011) *Mutation Research* 721, 163-170.
- Dertinger SD, Phonetheswath S, Avlasevich SL, Torous DK, Mereness J, Cottom J, Bemis JC, and MacGregor JT (2014) *Environmental & Molecular Mutagenesis* 55, 299-308.
- Dertinger SD, Phonetheswath S, Franklin D, Weller P, Torous DK, Bryce SM, Avlasevich S, Bemis JC, Hyrien O, Palis J, MacGregor JT (2010) *Toxicological Sciences* 115, 401-411.
- Dertinger SD, Heflich RH (2011) *Environmental Molecular Mutagenesis* 52, 681-684.
- Dertinger SD, Phonetheswath S, Avlasevich SL, Torous DK, Mereness J, Bryce SM, Bemis JC, Bell S, Weller P, MacGregor JT, (2012) *Toxicological Sciences* 130, 328-348.
- Dertinger SD, Phonetheswath S, Weller P, Avlasevich S, Torous DK, Mereness JA, Bryce SM, Bemis JC, Bell S, Portugal S, Aylott M, MacGregor JT (2011) *Environmental Molecular Mutagenesis* 52, 748-755.
- Dertinger SD, Phonetheswath S, Weller P, Nicolette J, Murray J, Sonders P, Vohr H-W, Shi J, Krsmanovic L, Gleason C, Custer L, Henwood A, Sweder K, Stankowski LF Jr., Roberts DJ, Giddings A, Kenny J, Lynch AM, Defrain C, Nesslany F, van der Leede B-j, Van Doninck T, Schuermans A, Tanaka K, Hiwata T, Tajima O, Wilde E, Elhajouji A, Gunther WC, Thiffeault CJ, Shutsky TJ, Fiedler RD, Kimoto T, Bhalli JA, Heflich RH, MacGregor JT (2011) *Environmental Molecular Mutagenesis* 52, 690-698.
- Dobo KL, Fiedler RD, Gunther WC, Thiffeault CJ, Cammerer Z, Coffing SL, Shutsky T, and Schuler M (2011) *Mutation Research* 725, 13-21.
- Dobrovolsky VN, Miura D, Heflich RH, Dertinger SD (2010) *Environmental Molecular Mutagenesis* 51, 825-835.
- Guérard M, Koenig J, Festag M, Dertinger SD, Singer T, Schmitt G, and Zeller A (2013) *Toxicological Sciences* 135, 309-316.
- Graupner A, Instanes C, Dertinger SD, Andersen JM, Lindeman B, Rongved TD, Brunborg G, and Olsen A (2014) *Mutation Research* 772, 34-41.
- Lemieux CL, Douglas GR, Gingerich J, Phonetheswath S, Torous DK, Dertinger SD, Phillips DH, Arlt VM, and White PA (2011) *Environmental & Molecular Mutagenesis* 52, 756-765.
- Lynch AM, Giddings A, Custer L, Gleason C, Henwood A, Aylott M, and Kenny J (2011) *Environmental & Molecular Mutagenesis* 52, 699-710.
- Patent numbers: 7,824,874, 8,062,860, 8,187,826, and patents pending.
- Phonetheswath S, Avlasevich SL, Torous DK, Mereness J, Bemis JC, Macgregor JT, and Dertinger SD (2013) *Environmental & Molecular Mutagenesis* 54, 294-298.
- Phonetheswath S, Franklin D, Torous DK, Bryce SM, Bemis JC, Raja S, Avlasevich S, Weller P, Hyrien O, Palis J, MacGregor JT, Dertinger SD (2010) *Toxicological Sciences* 114, 59-70.
- Schuler M, Gollapudi BB, Thybaud V, Kim JH (2011). *Environmental Molecular Mutagenesis* 52, 685-689.
- Shi J, Krsmanovic L, Bruce S, Kelly T, Paranjpe M, Szabo K, Arevalo M, Atta-Safah S, Debelie F, LaForce MK, Sly J, and Springer S (2011) *Environmental & Molecular Mutagenesis* 52, 711-720.
- Stankowski LF Jr, Roberts DJ, Chen H, Lawlor T, McKeon M, Murli H, Thakur A, and Xu Y (2011) *Environmental & Molecular Mutagenesis* 52, 738-747.
- Torous DK, Phonetheswath S, Avlasevich SL, Mereness J, Bryce SM, Bemis JC, Weller PC, Bell S, Gleason C, Custer LL, MacGregor JT, Dertinger SD (2012) *Environmental Molecular Mutagenesis* 53, 420-428.

## 10. Effective Date: August 01, 2021